

## FEATURED PHOTO

### RED-NAPED AND RED-BREASTED SAPSUCKERS

ROBERT A. HAMILTON, 34 Rivo Alto Canal, Long Beach, California 90803

JON L. DUNN, RR 2, Box 52R, Bishop, California 93514

The classification of the Yellow-bellied (*Sphyrapicus varius*), Red-naped (*S. nuchalis*), and Red-breasted (*S. ruber*) sapsuckers has confounded ornithologists since the late 1800s, a situation that spurred prodigious field and laboratory research during the past half-century aimed at elucidating relationships between these closely related taxa. The American Ornithologists' Union has recognized these three as species since 1985, citing the findings and conclusions of Johnson and Zink (1983). Here we consider relationships between *nuchalis* and *ruber*, sister species that are—intuition aside—more closely related to each other than either is to *varius* (Johnson and Zink 1983, Cicero and Johnson 1995). In addition to the scientific literature touched upon in this brief review, useful identification resources and overviews include Devillers (1970), DeBenedictis (1970), Dunn (1978), Kaufmann (1990), Lehman (1991), Pyle (1997), and Sibley (2000).

The Red-naped Sapsucker breeds in the Rocky Mountains, the eastern Cascades, and mountains of the Great Basin. The Red-breasted Sapsucker's northern subspecies, *S. r. ruber*, breeds from southeastern Alaska to southern Oregon, and apparently interbreeds extensively with southern *S. r. daggetti* in extreme southwestern Oregon and extreme northwestern California (Howell 1952). The breeding range of *S. r. daggetti* extends south to Marin County, across to northeastern California and south-central Oregon, and down through the Sierra Nevada, recurring in the high mountains of southern California (American Ornithologists' Union 1957, 1988).

The border between *S. r. ruber* and *S. nuchalis* traces the peaks of the Cascade Range north to central British Columbia, and mixing of these taxa "appears to be extremely scarce" (Howell 1952:268). Contact zones for *S. r. daggetti* and *S. nuchalis* have proven to be rather dynamic over the past century (Scott et al. 1976, Johnson and Johnson 1985), with two main zones of sympatry currently recognized. One encompasses parts of Modoc County, California, and adjacent Lake and Klamath counties, Oregon (Howell 1952). Examining 145 pairs in this area, Johnson and Johnson (1985) concluded that 18% of the studied pairs included at least one backcrossed individual, with interspecific breeding accounting for another 6%. The second zone of sympatry includes parts of eastern Mono County, California, and adjacent Lyon and Mineral counties, Nevada (Johnson and Johnson 1985), an area where interbreeding is "apparently very slight" (Howell 1952:251). At five additional known sites, *nuchalis* and *daggetti* are considered "essentially allopatric" by reason of one taxon being common and the other rare (Johnson and Johnson 1985). Otherwise, the Sierra Nevada is *daggetti* country, while *nuchalis* claims the Great Basin and Rockies; hybridization occurs in the areas of allopatry but is considered rare (Howell 1952; Johnson and Johnson 1985).

Both Red-breasted and Red-naped sapsuckers undergo prebasic molts on the breeding grounds. First-year Red-breasteds typically complete this molt by September, occasionally retaining some juvenal body feathers through December, and first-year Red-naped are about a month behind this schedule (Pyle 1997). Feathers of the lower throat appear to be the last to be replaced in both species (Howell 1952). Thus, birds seen in migration should be in fresh plumage (or mostly so). This is an important consideration, since the middle and basal portions of the head and chest feathers of all *daggetti* possess melanin pigment (Howell 1952), and wearing away of red feather tips normally reveals "an underlying 'shadow' that is reminiscent of the melanistic

## FEATURED PHOTO

breast band, cheek patch, and head stripes seen so vividly in *nuchalis*" (Johnson and Johnson 1985:4). The latter authors developed a 13-step "hybrid series index" that progresses from extreme *daggetti* (score 0) to extreme *nuchalis* (score 12), illustrating even "pure" *daggetti* with conspicuous black splotches on the upper breast; this apparently reflects the birds' appearance in summer, when body and head feathers are subject to considerable wear from attending to the nest and young. Note that Kaufman (1990) highlighted such blotches as an important feature of intermediate plumages, apparently because fall and winter Red-breasted Sapsuckers normally have unmarked red breasts.

Red-naped Sapsuckers winter in the southwestern U. S. from southern Nevada and southeastern California through central New Mexico, in western Mexico as far south as Jalisco, and throughout the Baja California peninsula (American Ornithologists' Union 1998). On the coastal slope of California, they are uncommon in San Diego County (Unitt 1984), become progressively rarer north to Santa Barbara County (Lehman 1994), and are very rare north of there to southern Oregon (American Ornithologists' Union 1998). Red-breasted Sapsuckers occupy a much smaller winter range along the Pacific slope from southeastern Alaska to northern Baja California, only occasionally straying into interior desert areas. Observers should be especially careful when identifying *daggetti* east of its normal range, as in Arizona, since many such claims involve hybrids (Rosenberg 1990, P. E. Lehman in litt.).

In mid-October 2001, while traversing White Pine and Lincoln counties in east-central Nevada, Dunn encountered approximately eight Red-naped Sapsuckers, four apparent hybrids, and no Red-breasted Sapsuckers. Observer coverage in this part of Nevada is light, but these numbers are consistent with Dunn's previous experience in the eastern deserts and the Great Basin. In the western and central parts of northern Baja California, where the normal ranges of Red-naped and Red-breasted sapsuckers overlap, the Red-naped is uncommon but outnumbered both the Red-breasted and hybrids—which are about equally rare—by ratios of about 5:1 (Erickson et al. 2001).

Dunn photographed the left-hand bird shown on the back cover of this issue on 13 October 2001 at Panamint Springs, Inyo County, California. Brian Small captured the other image during the month of October at Galileo Hill, Kern County, California. The birds' ages are uncertain, although both appear consistent with their first year. Neither is a "pure" Red-naped Sapsucker as both have too much red on their faces and show little or no trace of black frames on their throats. Unlike typical Red-breasted Sapsuckers, both birds have red pigment concentrated in the throat rather than spreading across the upper breast. Some dark feathering appears to be present below the right-hand bird's throat, with limited splotches of red below, as would be expected in an intermediate plumage, so this bird seems to be a fairly straightforward Red-naped × Red-breasted hybrid. Although the left-hand bird does not show obvious black splotching around the throat, its intermediate facial pattern and concentration of red in the throat, with only limited red below that, suggest another hybrid.

Thanks to Steve N. G. Howell and Kimball L. Garrett for assisting with the preparation of this note, and to Paul E. Lehman for his helpful review.

## LITERATURE CITED

- American Ornithologists' Union. 1957. Check-list of North American Birds, 5th ed. Am. Ornithol. Union, Washington, D. C.
- American Ornithologists' Union. 1985. Thirty-fifth supplement to the American Ornithologists' Union Check-list of North American Birds. Auk 102.
- American Ornithologists' Union. 1998. Check-list of North American Birds, 7th ed. Am. Ornithol. Union, Washington, D. C.

## FEATURED PHOTO

- Cicero, C., and Johnson, N. K. 1995. Speciation in Sapsuckers (*Sphyrapicus*): III. Mitochondrial DNA sequence divergence at the cytochrome-B locus. *Auk* 112:547–563.
- DeBenedictis, P. 1970. Gleanings from the technical literature. *Birding* 11:178–181.
- Devillers, P. 1970. Identification and distribution in California of the *Sphyrapicus varius* group of sapsuckers. *Calif. Birds* 1:47–76.
- Dunn, J. 1978. The races of the Yellow-bellied Sapsucker. *W. Tanager* 44:7.
- Erickson, R. A., Hamilton, R. A., and Howell, S. N. G. 2001. New information on migrant birds in northern and central portions of the Baja California Peninsula, including species new to Mexico, in *Birds of the Baja California Peninsula: Status, Distribution, and Taxonomy* (R. A. Erickson and S. N. G. Howell, eds.), pp. 112–170. Monogr. Field Ornithol. 3. Am. Birding Assoc., Colorado Springs, CO.
- Howell, T. R. 1952. Natural history and differentiation in the Yellow-bellied Sapsucker (*Sphyrapicus varius*). *Condor* 54:237–282.
- Kaufman, K. 1990. *A Field Guide to Advanced Birding*. Houghton Mifflin, Boston.
- Johnson, N. K., and Zink, R. M. 1983. Speciation in sapsuckers (*Sphyrapicus*): I. Genetic differentiation. *Auk* 100:871–884.
- Johnson, N. K., and Johnson, C. B. 1985. Speciation in sapsuckers (*Sphyrapicus*): II. Sympatry, hybridization, and mate preference in *Sphyrapicus ruber daggetti* and *S. nuchalis*. *Auk* 102:1–15.
- Lehman, P. 1991. Notes on plumage variation in adult Red-naped and Red-breasted sapsuckers. *Birding* 12:23–26.
- Lehman, P. E. 1994. *The Birds of Santa Barbara County, California*. Vert. Mus., Univ. Calif., Santa Barbara.
- Pyle, P. 1997. *Identification Guide to North American Birds, part I, Columbidae to Ploceidae*. Slate Creek Press, Bolinas, CA.
- Rosenberg, G. 1990. Arizona birding pitfalls, part 1: Species we take for granted. *Birding* 12:120–129.
- Scott, D. M., Ankey, C. D., and Jarosh, C. H. 1976. Sapsucker hybridization in British Columbia: Changes in 25 years. *Condor* 78:253–257.
- Sibley, D. A. 2000. *The Sibley Guide to Birds*. Knopf, New York.
- Unitt, P. 1984. The birds of San Diego County. *San Diego Soc. Nat. Hist. Mem.* 13.

# **27<sup>th</sup> Annual Meeting of the Western Field Ornithologists**

## **10–13 October 2002**

### **The San Joaquin Wildlife Sanctuary in Irvine and Ayres Country Inn in Costa Mesa, California**

#### Conference Information

Orange County, California, offers wonderful opportunities for birders to discover southern California resident and migrant bird species in rural, urban, suburban, and pelagic settings. The area is rich in bird life, and October will find WFO visiting during autumn migration. Bolsa Chica, Newport Back Bay, Limestone Canyon, Dripping Springs, the “gold coast,” the Pacific Ocean, Starr Ranch, and urban parrot roosts are examples of the field trips that have been planned for this meeting to highlight the rich diversity of landscape zones and birds that Orange County has to offer. With sufficient interest, postmeeting trips will go to the Salton Sea and Baja California.

#### Activities

WFO 2002 offers a mix of half-day field trips on October 11, 12, and 13. Other planned activities include a workshop on documenting bird records, evening field trips, a banding trip for hawks or owls, a full-day pelagic trip, afternoon paper sessions, and identification panels using both slides and recorded sounds. Social activities will include a welcoming reception, barbecue at the San Joaquin Wildlife Sanctuary, the annual dinner, and evening programs.

#### Speakers

The featured speakers for WFO 2002 are renowned marine biologist Robert L. Pitman and our own Philip Unitt. Bob will speak on the eastern Pacific pelagic frontier at the annual dinner to be held on Saturday, October 12. Phil will speak on the surprises revealed by the five years of field work for the San Diego County bird atlas at the barbecue on Friday, October 11.

#### Meeting Sponsors

WFO 2002 is locally hosted and sponsored by Sea and Sage Audubon Society and co-sponsored by Los Angeles Audubon Society. A number of local organizations are lending their support for this meeting.

#### Hotel Information

Headquarters for all conference activities will be Ayres Country Inn and Suites, 325 Bristol Street, Costa Mesa, California 92626. This is a world-class small hotel that is well located with easy airport or freeway access. Ask for Western Field Ornithologists when booking your reservations: 714-549-0300, 800-322-9992. Consider bringing the entire family. There are two pools and, for the enjoyment of nonbirding family members, free transportation to nearby sights and activities. The hotel has set aside a block of 50 double rooms for WFO at \$69.00 per night for a 3-night minimum stay, if the room is booked prior to September 15, 2002. The hotel is located 5 minutes from the Orange County/John Wayne Airport and will provide free transportation from the airport for hotel guests. The conference schedule, costs, and registration forms are posted at [www.wfo-cbrc.org](http://www.wfo-cbrc.org) and [www.seaandsageaudubon.org](http://www.seaandsageaudubon.org).

For additional information contact [robcatwaters@earthlink.net](mailto:robcatwaters@earthlink.net).

# WESTERN BIRDS

World Wide Web site:  
[www.wfo-cbrc.org](http://www.wfo-cbrc.org)

Quarterly Journal of Western Field Ornithologists

*President:* Mike San Miguel, 2132 Highland Oaks Dr., Arcadia, CA 91006;  
sanmigbird@aol.com

*Vice-President:* Daniel D. Gibson, University of Alaska Museum, 907 Yukon Dr.,  
Fairbanks, AK 99775-6960

*Treasurer/Membership Secretary:* Robbie Fischer, 1359 Solano Drive, Pacifica, CA  
94044; robbie22@pacbell.net

*Recording Secretary:* Kei Sochi, The Nature Conservancy, Western Regional Of-  
fice, 2060 Broadway, Suite 230, Boulder, CO 80302; ksochi@tnc.org

*Directors:* Ted Floyd, Kimball L. Garrett, Daniel D. Gibson, Bob Gill, Gjon Hazard,  
Dave Krueper, Mike San Miguel, W. David Shuford, David Yee

*Editor:* Philip Unitt, San Diego Natural History Museum, P. O. Box 121390, San  
Diego, CA 92112-1390; birds@sdnhm.org

*Associate Editors:* Daniel D. Gibson, Robert A. Hamilton, Ronald R. LeValley,  
Tim Manolis, Kathy Molina, Mark K. Sogge

*Graphics Manager:* Virginia P. Johnson, 4637 Del Mar Ave., San Diego, CA 92107

*Photo Editor:* Peter La Tourrette, 1019 Loma Prieta Ct., Los Altos, CA 92024

*Featured Photo:* Robert A. Hamilton, 34 Rivo Alto Canal, Long Beach, CA 90803

*Book Reviews:* Steve N. G. Howell, Point Reyes Bird Observatory, 4990 Shoreline  
Highway, Stinson Beach, CA 94970

*Secretary, California Bird Records Committee:* Guy McCaskie, P. O. Box 275,  
Imperial Beach, CA 91933-0275; guymcc@pacbell.net

*Chairman, California Bird Records Committee:* Richard A. Erickson, LSA Associates,  
1 Park Plaza, Suite 500, Irvine, CA 92614; richard.erickson@lsa-assoc.com

Membership dues, for individuals and institutions, including subscription to *Western Birds*: Patron, \$1000.00; Life, \$400 (payable in four equal annual installments); Supporting, \$60 annually; Contributing, \$34 annually; Family \$26; Regular U.S. \$22 for one year, \$41 for two years, \$60 for three years. Dues and contributions are tax-deductible to the extent allowed by law.

Send membership dues, changes of address, correspondence regarding missing issues, and orders for back issues and special publications to the Treasurer. Make checks payable to Western Field Ornithologists.

Back issues of *Western Birds* within U.S. \$24 per volume, \$6.00 for single issues, plus \$1.00 for postage. Outside the U.S. \$30 per volume, \$7.50 for single issues.

The California Bird Records Committee of Western Field Ornithologists revised its 10-column Field List of California Birds in January 2000. The last list covered 606 accepted species; the new list covers 613 species. Please send orders to WFO, c/o Robbie Fischer, Treasurer, 1359 Solano Drive, Pacifica, CA 94044. California addresses please add 7.75% sales tax.

Quantity: 1-9, \$1.50 each, includes shipping and handling; 10-39, \$1.30 each, add \$2.00 for shipping and handling; 40 or more, \$1.15 each, add \$4.00 for shipping and handling.